The **Juvenile Arthritis Disease Activity Score (JADAS)** is composed of the following measures: 1) physician's global assessment (0–10); 2) parent's global assessment (0–10); 3) active joint count (assessed in 71, 27 or 10 joints); 4) ESR (normalized to 0–10).

	JADAS-71	JADAS-27	JADAS-10
Physician global assessment	0–10-cm VAS	0–10-cm VAS	0–10-cm VAS
Parent/patient global assessment	0–10-cm VAS	0–10-cm VAS	0–10-cm VAS
Active joint count	Simple, 0–71 joints	Simple, 0–27 joints	Simple, 0–10 joints†
Swollen joint count (range)	–	–	–
Tender joint count (range)	–	–	–
Acute-phase reactant (range)	Normalized ESR (0–10)‡	Normalized ESR (0–10)‡	Normalized ESR (0–10)‡
Score range	0–101	0–57	0–40

* JADAS-71 = Juvenile Arthritis Disease Activity Score in 71 joints; DAS28 = Disease Activity Score in 28 joints; CDAI = Clinical Disease Activity Index; VAS = visual analog scale; ESR = erythrocyte sedimentation rate.
† Up to 10 joints, irrespective of their type, censored at 10.
‡ According to the formula: (value in mm/hour −20)/10, where values <20 mm/hour are converted to 0, and values >120 mm/hour are converted to 120.

JADAS-71



JADAS-27



References:

Consolaro, A., Giancane, G., Schiappapietra, B. et al. Clinical outcome measures in juvenile idiopathic arthritis. Pediatr Rheumatol 14, 23 (2016). https://doi.org/10.1186/s12969-016-0085-5

Consolaro A, Ruperto N, Bazso A, Pistorio A, Magni-Manzoni S, Filocamo G, et al. Development and validation of a composite disease activity score for juvenile idiopathic arthritis. Arthritis Rheum. 2009;61:658–66.

Jeong, D.C. (2014). Assessment of Disease Activity in Juvenile Idiopathic Arthritis. Journal of Reproduction and Development, 21, 289-296.